

# Public Health Outcomes Framework: Update for Leicestershire

August 2018

**Janine Johnson**

Public Health Analyst

Strategic Business Intelligence Team  
Strategy Business Intelligence Branch  
Chief Executive's Department  
Leicestershire County Council  
County Hall, Glenfield  
Leicester LE3 8RA

Tel 0116 305 4257

Email [janine.johnson@leics.gov.uk](mailto:janine.johnson@leics.gov.uk)


Produced by the Strategic Business Intelligence Team at Leicestershire County Council.

Whilst every effort has been made to ensure the accuracy of the information contained within this report, Leicestershire County Council cannot be held responsible for any errors or omission relating to the data contained within the report.

# What have we learnt in the PHOF update?

## 1. First time entrants to the youth justice system continue to perform well


- The recent trend for first time entrants to the youth justice system shows Leicestershire and Rutland (combined) have significantly declined and have performed significantly better than England average for the last five years.
- Compared to the previous year, the rate of first time entrants to the youth justice system in Leicestershire and Rutland has increased from 166.7 per 100,000 population aged 10-17 years in 2016 to 180.5 per 100,000 population aged 10-17 years in 2017, this equates to an increase of 9 first time entrants to the youth justice system.
- This increase is not reflected nationally as the England rate continues to decrease from 331.0 per 100,000 population aged 10-17 years in 2016 to 292.5 per 100,000 population aged 10-17 years in 2017.



## 2. Deaths from drug misuse continues to rise

### Deaths from drug misuse


- The rate of deaths from drug misuse has increased year on year for the last three reporting periods both in Leicestershire and nationally. Despite this, Leicestershire has continued to perform significantly better than the national average since this indicator was recorded.
- Compared to the previous year, the number of deaths from drug misuse has increased from 33 in 2014-16 to 42 in 2015-17 reflecting an increase of 9 deaths.
- In 2015-17, the rate of deaths from drug misuse is 2.2 per 100,000 population in Leicestershire. This is half the national rate of 4.3 per 100,000 population in 2015-17.



## 3. Smoking in adults is on the decrease

### Smoking prevalence in adults

- In 2017 in Leicestershire, 12.1% of adults aged 18+ smoked. This is significantly better than the England average of 14.9%. This is the first year (2017) Leicestershire has performed significantly better than England.
- Compared to the previous year, the number of current smokers has decreased from 73,316 in 2016 to 65,688 adults in 2017. This is a decrease of 7,628 adult smokers.
- In 2017 in Leicestershire, the smoking prevalence between the genders is similar; this pattern is not witnessed nationally. In Leicestershire the prevalence for males is 12.1% and for females is 12.0%. This is similar to the national average for females (13.0%) but significantly lower than the national average for males (16.8%).



- In 2017, the smoking prevalence of working age adults (18-64 years) in Leicestershire is 13.6%. Adults who work in a managerial role have a significantly better prevalence of smoking prevalence (10.0%) compared to the Leicestershire average. Adults who work in a routine role have a smoking prevalence of 18.7%, this is the highest prevalence out of all socioeconomic groups but is still similar to the Leicestershire average.

#### 4. Health checks – data quality issue

##### Cumulative percentage of the eligible population aged 40-74 who were offered an NHS health check

- There is a data quality issue with Leicestershire’s value: This is because 240,560 people were offered an NHS health check whereas the total eligible population is 206,998. This means over 100% (116.2%) were offered an NHS health check between 2013/14 – 2017/18. This is why the PHOF states the value for this indicator is 100%.
- Over the same period, nationally 90.9% of the eligible population aged 40-74 was offered an NHS health check.

##### Cumulative percentage of the eligible population aged 40-74 who were offered and received an NHS health check

- Over a cumulative five year period (2013/14-2017/18), 43.1% of the eligible population of Leicestershire aged 40-74 years who were offered an NHS health check received an NHS health check. This is calculated as 103,772 people aged 40-74 eligible for an NHS Health Check who have received an NHS Health Check in the five year period divided by 240,560 people aged 40-74 eligible for an NHS Health Check who were offered an NHS Health Check in the five year period. This local percentage is significantly worse than the national average (48.7%).


##### Cumulative percentage of the eligible population aged 40-74 who received an NHS health check

- Over a cumulative five year period (2013/14-2017/18), 50.1% of the eligible population of Leicestershire aged 40-74 years received an NHS health check. This is calculated as 103,772 people aged 40-74 eligible for an NHS Health Check who received an NHS Health Check in the five year period divided by 206,998 people aged 40-74 eligible for an NHS Health Check in the five year period. Leicestershire performs significantly better than the national average of 44.3%.

#### 5. Flu vaccination coverage continues to perform badly compared to the benchmark


##### Population vaccination coverage – flu (aged 65+) (*benchmark goal of higher than 75%*)

- For eight consecutive years Leicestershire and Rutland (combined) have performed worse against the benchmark goal of higher than 75%. This pattern is also reflected nationally.
- Despite this, Leicestershire and Rutland have performed significantly higher than the national average since 2013/14. The local trend has also shown a significant increase over the last eight time periods.
- In 2017/18, 112,024 eligible adults aged 65+ in Leicestershire and Rutland received the flu vaccine, this equates to a coverage percentage of 74.1%. This is significantly higher than the national average of 72.6%.



**Population vaccination coverage – flu (at risk individuals) (benchmark goal of higher than 55%)**

- For eight consecutive years Leicestershire and Rutland (combined) have performed worse than the benchmark goal of higher than 55%. This pattern is also reflected nationally.
- The local trend has shown a significant decline over the last five time periods.
- In 2017/18, 39,668 at risk individuals aged 6 months to under 65 years (excluding healthy pregnant women and carers) in Leicestershire and Rutland received the flu vaccine, this equates to a coverage percentage of 48.5%. The national coverage is 48.9%.




**Population vaccination coverage – flu (2-3 year olds) – new definition**

- The age group for indicator 3.03xviii (Flu vaccination aged 2-3 years old) has been changed from 2-4 years old as 4 year olds are no longer vaccinated in a GP setting.
- In 2017/18, 61.4% of children aged 2-3 years in Leicestershire and Rutland received the flu vaccine, this is similar to the benchmarking goal of 40% - 65%.

**6. Significant increase in children free from tooth decay in five year olds**

**Proportion of five year old children free from dental decay**


- Of the five year olds in Leicestershire who were examined for tooth decay in 2016/17, 77.7% were reported as being free from obvious dental decay. This equates to 2,228 children free from dental decay, more than double the amount of children (1,106) reported in 2014/15.
- In 2016/17 Leicestershire performed similar to the national average of 76.7%. Previously, since 2007/08 Leicestershire had been significantly worse than England. Between 2014/15 and 2016/17, a significant increase in the proportion of five year old children free from dental decay was witnessed.



**7. Chlamydia detection rate is worse than the benchmark goal**

**Chlamydia detection (benchmark goal of 1,900 per 100,000 population aged 15-24 years)**

- In Leicestershire the chlamydia detection rate decreased (got worse) from a rate of 1,973 per 100,000 population aged 15-24 years in 2016 to 1,887 per 100,000 population aged 15-24 years in 2017. This equates to a decrease of 74 diagnoses.
- The rate has declined to be worse than the benchmark of <1,900 in 2017, whereas in 2016 the county performed similar to the benchmark goal
- A decline in rate between 2016 and 2017 was witnessed in both males and females in Leicestershire.




To note, the national rate of 1,882 per 100,000 is now rated significantly worse against the benchmark goal of 1,900 per 100,000 population aged 15-24 years.

## 8. Estimated dementia diagnosis rate remains similar to the benchmark

### Estimated dementia diagnosis rate (aged 65+) (benchmark goal of 66.7%)

- In 2018, 6,089 Leicestershire residents aged 65+ who were registered with a GP had a diagnosis of dementia. This equates to 70.0% of the estimated number (8,695) of Leicestershire residents aged 65+ whom would have been expected to have dementia have been diagnosed.



# Appendix 1: Performance Summary


This release of the PHOF contained 253 indicators (including gender breakdowns). The overall findings are summarised below.

Overall performance across Leicestershire County is significantly better than the England average or against a benchmarking goal for 114 of the indicators presented in the PHOF. Performance is significantly worse for 18 indicators and these indicators need to show improvement.

**Table 1: Summary of RAG Ratings for Leicestershire August 2018**

Domain	Better	Similar	Worse	Not compared	Total
Overarching	8			8	16
Domain 1: Wider determinants of health	21	20	10	10	61
Domain 2: Health improvement	39	21	4	14	78
Domain 3: Health protection	13	6	3	8	30
Domain 4: Healthcare premature mortality	33	33	1	1	68
<b>Total</b>	<b>114</b>	<b>80</b>	<b>18</b>	<b>41</b>	<b>253</b>

**Figure 1: Count of Leicestershire PHOF RAG ratings, August 2018**



## 1. All Indicators - Leicestershire Performance Against Nearest Neighbours

Nearest CIPFA neighbours to Leicestershire available from PHOF tool include:

- Cambridgeshire
- Derbyshire
- Devon
- Essex
- Gloucestershire
- Hampshire
- North Yorkshire
- Northamptonshire
- Nottinghamshire
- Oxfordshire
- Somerset
- Staffordshire
- Suffolk
- Warwickshire
- Worcestershire

Leicestershire's performance for the updated indicators ranked in relation to nearest neighbours is summarised in Appendix 2. Best or lowest\* performance is ranked 1. The value for each indicator for Leicestershire along with best and worse cluster performance is also provided.

*\* when no polarity has been applied, the indicator is ranked from low to high.*

### Poor Performing Indicators Against Nearest Neighbours

There are currently 21 indicators where, when compared to similar areas, Leicestershire performs in the bottom three (worse performing). These are:

#### Domain 1: Wider Determinants of Health

- 1.02i - School Readiness: % of children with FSM status achieving a GLOD at reception 2016/17 (P) 16/16
- 1.02ii - School Readiness: % of Year 1 pupils with FSM status achieving level in the phonics check 2016/17 (P) 16/16
- 1.08iii - Gap in employment for those in contact with sec MH services and overall rate 2016/17 (P) 16/16
- 1.08iii - Gap in employment for those in contact with sec MH services and overall rate 2016/17 (F) 16/16

#### Domain 2: Health Improvement

- 2.05ii – Proportion of children aged 2-2½yrs offered ASQ-3 as part of the HCP or integrated review 2016/17 (P) 16/16
- 2.08i - Average difficulties score for all LAC aged 5-16 who have been in care => 12 months 2016/17 (P) 15/16
- 2.08ii - Percentage of children where there is a cause for concern 2016/17 (P) 14/16
- 2.11iii - Average number of portions of vegetables consumed daily (adults) 2016/17 (P) 15/16
- 2.20iv - Abdominal Aortic Aneurysm Screening - Coverage 2016/17 (M) 15/16



### **Domain 3: Health Protection**

- 3.01 - Fraction of mortality attributable to particulate air pollution 2016 (P) 16/16
- 3.06 – NHS organisations with a board approved sustainable development management plan 2015/16 (n/a) 15/16

### **Domain 4: Healthcare Public Health and Preventing Premature Mortality**

- 4.02 - Proportion of five year old children free from dental decay 2016/17 (P) 15/16
- 4.11 - Emergency readmissions within 30 days of discharge from hospital 2011/12 (F) 14/16
- 4.11 - Emergency readmissions within 30 days of discharge from hospital 2011/12 (P) 14/16
- 4.14i - Hip fractures in people aged 65 and over 2016/17 (F) 15/15
- 4.14i - Hip fractures in people aged 65 and over 2016/17 (P) 15/15
- 4.14ii - Hip fractures in people aged 65 and over - aged 65-79 2016/17 (F) 13/15
- 4.14ii - Hip fractures in people aged 65 and over - aged 65-79 2016/17 (P) 13/15
- 4.14iii - Hip fractures in people aged 65 and over - aged 80+ 2016/17 (F) 15/15
- 4.14iii - Hip fractures in people aged 65 and over - aged 80+ 2016/17 (P) 13/15
- 4.15ii – Excess winter deaths index (single year, aged 85+) Aug 2015 – July 2016 (M) 14/16

## **2. All Indicators - Leicestershire's Performance is Getting Worse**

From August 2016, markers of significant trend were added to the PHOF. When there are at least five consecutive non-overlapping data points available for a proportion or crude rate indicator, a chi-squared statistical test for trend is carried out on the most recent five points. If there is no significant trend in the most recent five points, the test is redone, including an additional (sixth) point, and so on, going further back until the full series has been tested.

The following indicators have a most recent trend of “getting worse” in Leicestershire:

- 2.07ii - Hospital admissions caused by unintentional and deliberate injuries in young people (aged 15-24 years) 2016/17 (P)
- 2.15i - Successful completion of drug treatment - opiate users 2016 (P)
- 2.20i - Cancer screening coverage - breast cancer 2017 (F)
- 2.20ii - Cancer screening coverage - cervical cancer 2017 (F)
- 3.03vi - Population vaccination coverage - Hib / MenC booster (2 years old) 2016/17 (P)
- 3.03xv - Population vaccination coverage - Flu (at risk individuals) 2017/18 (P)

## **3. All Indicators - Leicestershire Performance against National**

Currently, performance is significantly worse for 18 indicators and these indicators need to show improvement. These are:

### **Domain 1: Wider Determinants of Health**

- 1.02i - School Readiness: % of children with free school meal status achieving a good level of development at reception 2016/17 (F)
- 1.02i - School Readiness: % of children with free school meal status achieving a good level of development at reception 2016/17 (P)
- 1.02ii - School Readiness: % of Year 1 pupils with free school meal status achieving the expected level in the phonics check 2016/17 (F)

- 1.02ii - School Readiness: % of Year 1 pupils with free school meal status achieving the expected level in the phonics check 2016/17 (M)
- 1.02ii - School Readiness: % of Year 1 pupils with free school meal status achieving the expected level in the phonics check 2016/17 (P)
- 1.06ii - Adults with secondary Mental Health services who live in stable and appropriate accommodation, 2016/17 (F)
- 1.06ii - Adults with secondary Mental Health services who live in stable and appropriate accommodation, 2016/17 (M)
- 1.06ii - Adults with secondary Mental Health services who live in stable and appropriate accommodation, 2016/17 (P)
- 1.08iii - Gap in employment for those in contact with secondary Mental Health services and overall employment rate 2016/17 (F)
- 1.08iii - Gap in employment for those in contact with secondary Mental Health services and overall employment rate 2016/17 (P)

#### **Domain 2: Health Improvement**

- 2.05ii - Prop of children aged 2-2½yrs offered ASQ-3 as part of the HCP or integrated review 2016/17 (P)
- 2.08ii - Percentage of children where there is a cause for concern 2015/16 (P)
- 2.11iii – Average number of portions of vegetables consumed daily 2016/17 (P)
- 2.22iv- Cumulative % of eligible pop aged 40-74 offered and received an NHS Health Check 2013/14 - 16/17 (P)

#### **Domain 3: Health Protection**

- 3.02 – Chlamydia detection rate (15-24 years) 2017 (P)
- 3.03xiv – Population vaccination coverage – Flu (aged 65+) 2017/18 (P)
- 3.03xv – Population vaccination coverage – Flu (at risk individuals) 2017/18 (P)

#### **Domain 4: Healthcare Public Health and Preventing Premature Mortality**

- 4.12iv - Preventable sight loss - sight loss certifications 2015/16 (P)

# Overarching Indicators: Leicestershire

Indicator	Value	Rank	Best/Lowest	Worst/Highest	England	DoT	RAG
0.1i - Healthy life expectancy at birth, (F), 2014 - 16	65.8	8/16	68.7	62.4	63.9		●
0.1i - Healthy life expectancy at birth, (M), 2014 - 16	65.2	8/16	67.5	61.7	63.3		●
0.1ii - Life expectancy at 65, (F), 2014 - 16	21.7	8/16	22.0	20.7	21.1		●
0.1ii - Life expectancy at 65, (M), 2014 - 16	19.3	7/16	19.8	18.4	18.8		●
0.1ii - Life expectancy at birth, (F), 2014 - 16	84.0	8/16	84.6	82.7	83.1		●
0.1ii - Life expectancy at birth, (M), 2014 - 16	80.7	5/16	81.4	79.1	79.5		●
0.2i - Inequality in life expectancy at birth ENGLAND, (F), 2014 - 16	Null	Null	Null	Null	7.3		●
0.2i - Inequality in life expectancy at birth ENGLAND, (M), 2014 - 16	Null	Null	Null	Null	9.3		●
0.2iii - Inequality in life expectancy at 65 LA, (F), 2014 - 16	3.5	11/16	2.1	5.3	Null		●
0.2iii - Inequality in life expectancy at 65 LA, (M), 2014 - 16	3.5	9/16	2.5	5.0	Null		●
0.2iii - Inequality in life expectancy at birth LA, (F), 2014 - 16	5.3	10/16	4.0	8.1	Null		●
0.2iii - Inequality in life expectancy at birth LA, (M), 2014 - 16	6.2	5/16	4.9	9.4	Null		●
0.2iv - Gap in life expectancy at birth between each local authority and England as a whole, (F), 2014 - 16	0.9	8/16	1.4	-0.4	0.0		●
0.2iv - Gap in life expectancy at birth between each local authority and England as a whole, (M), 2014 - 16	1.2	5/16	1.8	-0.4	0.0		●
0.2vi - Inequality in healthy life expectancy at birth LA, (F), 2009 - 13	9.6	5/16	7.1	14.4	Null		●
0.2vi - Inequality in healthy life expectancy at birth LA, (M), 2009 - 13	9.0	3/16	8.3	14.9	Null		●

**Statistical Significance compared to England or Benchmark:**  
 ● Better ● Same ● Worse  
 ● Higher ● Lower ● Not compared  
**Direction of Travel:**  
 ▲ Increasing ▼ Decreasing  
 ▲ Increasing and getting better ▼ Decreasing and getting better  
 ▲ Increasing and getting worse ▼ Decreasing and getting worse  
 ▲ No significant change ▼ Cannot be calculated

Ranking, Best and Worst columns are compared to the nearest neighbours only. Rank: 1 is calculated as the best (or lowest when no polarity is applied)

Source: August 2018 PHOF, PHE

# Wider Determinants of Health: Leicestershire

Indicator	Value	Rank	Best/Lowest	Worst/Highest	England	DoT	RAG
1.01i - Children in low income families (all dependent children under 20), (P), 2015	10.1	4/16	9.6	14.7	16.6	▼	●
1.01ii - Children in low income families (under 16s), (P), 2015	10.4	4/16	9.8	15.1	16.8	▼	●
1.02i - School Readiness: % of children achieving a GLOD at reception, (F), 2016/17	76.6	4/16	74.9	82.6	77.7	▲	●
1.02i - School Readiness: % of children achieving a GLOD at reception, (M), 2016/17	64.1	8/16	60.6	69.3	64.0	▲	●
1.02i - School Readiness: % of children achieving a GLOD at reception, (P), 2016/17	70.1	12/16	75.7	68.1	70.7	▲	●
1.02i - School Readiness: % of children with FSM status achieving a GLOD at reception, (F), 2016/17	50.4	1/16	50.4	70.8	64.4	▲	●
1.02i - School Readiness: % of children with FSM status achieving a GLOD at reception, (M), 2016/17	43.0	6/16	38.4	51.5	48.1	▲	●
1.02i - School Readiness: % of children with FSM status achieving a GLOD at reception, (P), 2016/17	46.4	16/16	59.3	46.4	56.0	▲	●
1.02ii - School Readiness: % of Yr 1 pupils achieving the level in phonics check, (F), 2016/17	84.6	8/16	82.4	86.9	84.8	▲	●
1.02ii - School Readiness: % of Yr 1 pupils achieving the level in phonics check, (M), 2016/17	78.7	12/16	75.0	79.6	77.6	▲	●
1.02ii - School Readiness: % of Yr 1 pupils achieving the level in phonics check, (P), 2016/17	81.6	6/16	83.1	78.8	81.1	▲	●
1.02ii - School Readiness: % of Yr 1 pupils with FSM status ach level in the phonics check, (F), 2016/17	63.5	1/16	63.5	76.3	73.9	▲	●
1.02ii - School Readiness: % of Yr 1 pupils with FSM status ach level in the phonics check, (M), 2016/17	56.6	5/16	52.4	65.6	63.2	▲	●
1.02ii - School Readiness: % of Yr 1 pupils with FSM status ach level in the phonics check, (P), 2016/17	59.6	16/16	69.0	59.6	68.4	▲	●
1.03 - Pupil absence, (P), 2016/17	4.4	4/16	4.3	4.9	4.7	▼	●
1.04 - First time entrants to the youth justice system, (P), 2017	180.5	4/16	131.6	352.3	292.5	▼	●
1.05 - 16-17 year olds NEET or whose activity is not known - current method, (F), 2016	4.7	6/16	3.3	7.1	5.4	—	●
1.05 - 16-17 year olds NEET or whose activity is not known - current method, (M), 2016	5.4	6/16	4.1	8.5	6.6	—	●
1.05 - 16-17 year olds NEET or whose activity is not known - current method, (P), 2016	5.1	6/16	3.7	7.5	6.0	—	●
1.05 - 16-18 year olds NEET - historical method, (P), 2015	3.0	3/16	2.5	5.2	4.2	▼	●
1.06i - Adults with a learning disability who live in stable and appropriate accom, (F), 2016/17	79.4	10/16	66.9	89.4	76.8	▲	●
1.06i - Adults with a learning disability who live in stable and appropriate accom, (M), 2016/17	79.4	12/16	64.5	88.6	75.8	▲	●
1.06i - Adults with a learning disability who live in stable and appropriate accom, (P), 2016/17	79.4	5/16	88.9	65.8	76.2	▲	●
1.06ii - Adults with secondary MH services who live in stable and appropriate accom., (F), 2016/17	37.0	3/16	22.0	88.0	56.0	—	●
1.06ii - Adults with secondary MH services who live in stable and appropriate accom., (M), 2016/17	37.0	5/16	26.0	84.0	53.0	—	●
1.06ii - Adults with secondary MH services who live in stable and appropriate accom., (P), 2016/17	37.0	13/16	86.0	24.0	54.0	—	●
1.08i - Gap in the employment rate between those with LTC and the overall rate, (P), 2016/17	30.9	2/16	27.6	37.6	29.4	—	●
1.08ii - Gap in the employment rate between those with LD and the overall rate, (F), 2016/17	66.9	7/16	55.2	74.1	64.5	—	●
1.08ii - Gap in the employment rate between those with LD and the overall rate, (M), 2016/17	66.7	1/16	66.7	81.6	73.3	—	●
1.08ii - Gap in the employment rate between those with LD and the overall rate, (P), 2016/17	66.3	2/16	65.3	77.1	68.7	—	●
1.08iii - Gap in employment for those in contact with sec MH services and overall rate, (F), 2016/17	71.3	16/16	50.0	71.3	60.5	—	●
1.08iii - Gap in employment for those in contact with sec MH services and overall rate, (M), 2016/17	76.7	13/16	66.8	79.5	73.5	—	●
1.08iii - Gap in employment for those in contact with sec MH services and overall rate, (P), 2016/17	74.4	16/16	59.3	74.4	67.4	—	●
1.08iv - Percentage of people aged 16-64 in employment, (F), 2016/17	75.3	13/16	69.0	77.6	69.5	▲	●
1.08iv - Percentage of people aged 16-64 in employment, (M), 2016/17	79.7	3/16	78.2	85.5	79.5	▲	●
1.08iv - Percentage of people aged 16-64 in employment, (P), 2016/17	77.4	9/16	80.8	74.7	74.4	▲	●

**Statistical Significance compared to England or Benchmark:**  
● Better ● Same ● Worse  
▲ Higher ● Lower ● Not compared  
▼ Decreasing ▲ Increasing ▲ No significant change  
▼ Decreasing and getting better ▲ Increasing and getting better — Cannot be calculated  
▼ Decreasing and getting worse ▲ Increasing and getting worse

Ranking, Best and Worst columns are compared to the nearest neighbours only. Rank: 1 is calculated as the best (or lowest when no polarity is applied)

Source: August 2018 PHOF, PHE

## Wider Determinants of Health (cont): Leicestershire

Indicator	Value	Rank	Best/Lowest	Worst/Highest	England	DoT	RAG
1.09i - Sickness absence - % who had at least one day off in the previous week, (P), 2014 - 16	1.8	4/16	1.6	2.9	2.1	■	●
1.09ii - Sickness absence - % of working days lost due to sickness absence, (P), 2014 - 16	1.0	5/16	0.7	1.8	1.2	■	●
1.10 - Killed and seriously injured (KSI) casualties on England's roads, (P), 2014 - 16	35.4	4/16	28.1	71.3	39.7	■	●
1.11 - Domestic abuse - historic method, (P), 2014/15	20.6	12/16	13.0	22.7	20.4	■	●
1.11 - Domestic abuse-related incidents and crimes - current method, (P), 2016/17	18.7	6/16	14.8	28.5	22.5	■	●
1.12i - Violent crime (including sexual violence) - hospital admissions for violence, (F), 2014/15 - 16/17	9.1	4/15	6.1	18.2	17.2	■	●
1.12i - Violent crime (including sexual violence) - hospital admissions for violence, (M), 2014/15 - 16/17	34.9	2/15	18.9	70.9	68.4	■	●
1.12i - Violent crime (including sexual violence) - hospital admissions for violence, (P), 2014/15 - 16/17	22.1	3/15	12.5	44.6	42.9	■	●
1.12ii - Violent crime (inc sexual violence) - violence offences per 1,000 pop, (P), 2016/17	11.1	3/16	9.5	21.9	20.0	▲	●
1.12iii - Violent crime (inc sexual violence) - rate of sexual offences per 1,000 pop, (P), 2016/17	1.1	1/16	1.1	2.4	1.9	▲	●
1.13i - Re-offending levels - percentage of offenders who re-offend, (P), 2014	20.7	1/16	20.7	26.7	25.4	▲	●
1.13ii - Re-offending levels - average number of re-offences per offender, (P), 2014	0.6	2/16	0.6	0.9	0.8	▲	●
1.13iii - First time offenders, (P), 2017	117.0	6/16	71.3	205.2	166.4	▼	●
1.14i - The rate of complaints about noise, (P), 2015/16	3.2	3/16	2.6	5.3	6.3	▲	●
1.14ii - % exposed to road, rail, air transport noise of 65dB(A) or more, during the day, (P), 2011	2.5	2/16	2.5	4.0	5.2	■	●
1.14iii - % exposed to road, rail, air transport noise of 55 dB(A) or more during the night, (P), 2011	5.1	9/16	3.8	7.5	8.0	■	●
1.15i - Statutory homelessness - Eligible homeless people not in priority need, (P), 2016/17	0.4	10/16	0.2	1.1	0.8	■	●
1.15ii - Statutory homelessness - households in temporary accommodation, (P), 2016/17	0.4	5/16	0.2	2.8	3.3	■	●
1.16 - Utilisation of outdoor space for exercise/health reasons, (P), Mar 2015 - Feb 2016	20.8	3/16	24.4	14.2	17.9	■	●
1.17 - Fuel poverty, (P), 2015	11.4	8/16	6.8	12.3	11.0	▲	●
1.18i - Social Isolation: % of SC users with as much social contact as they would like, (P), 2016/17	46.2	9/16	52.8	42.0	45.4	■	●
1.18ii - Social Isolation: % of adult carers with as much social contact as they would like, (P), 2016/17	31.4	11/16	45.4	24.9	35.5	■	●

**Statistical Significance compared to England or Benchmark:**

- Better
- Same
- Worse
- Higher
- Lower
- Not compared

**Direction of Travel:**

- ▲ Increasing
- ▼ Decreasing
- ▲ Increasing and getting better
- ▼ Decreasing and getting better
- ▲ Increasing and getting worse
- ▼ Decreasing and getting worse

▲ No significant change  
 ▼ Cannot be calculated

Ranking, Best and Worst columns are compared to the nearest neighbours only. Rank: 1 is calculated as the best (or lowest when no polarity is applied)

Source: August 2018 PHOF, PHE

# Health Improvement: Leicestershire

Indicator	Value	Rank	Best/Lowest	Worst/Highest	England	DoT	RAG
2.01i - Low birth weight of term babies, (P), 2016	2.4	10/16	1.8	2.7	2.8	▲	●
2.02i - Breastfeeding initiation, (F), 2016/17	Null	Null	Null	Null	74.5	■	●
2.02ii - Breastfeeding prevalence at 6-8 weeks after birth - current method, (P), 2016/17	Null	Null	Null	Null	44.4	■	●
2.03 - Smoking status at time of delivery - current method, (F), 2016/17	8.6	2/16	7.7	14.8	10.7	▼	●
2.03 - Smoking status at time of delivery - historical method, (F), 2016/17	8.6	3/14	7.7	14.5	10.5	▼	●
2.04 - Under 18 conceptions, (F), 2016	13.7	5/16	11.6	22.4	18.8	▼	●
2.04 - Under 18 conceptions: conceptions in those aged under 16, (F), 2016	2.1	6/16	1.4	3.6	3.0	▼	●
.. children aged 2-2½yrs receiving ASQ-3 as part of the Healthy Child Programme or integrated review, (P), 2016/17	41.6	16/16	100.0	41.6	89.4	■	●
2.06i - Child excess weight in 4-5 and 10-11 year olds - 4-5 year olds, (P), 2016/17	20.3	3/16	18.5	24.9	22.6	▲	●
2.06ii - Child excess weight in 4-5 and 10-11 year olds - 10-11 year olds, (P), 2016/17	29.6	4/16	27.1	34.2	34.2	▲	●
2.07i - Hosp adm caused by unintentional and deliberate injuries in children (aged 0-4 yrs), (F), 2016/17	74.4	1/15	74.4	146.9	112.0	■	●
2.07i - Hosp adm caused by unintentional and deliberate injuries in children (aged 0-4 yrs), (M), 2016/17	86.5	1/15	86.5	183.1	140.0	■	●
2.07i - Hosp adm caused by unintentional and deliberate injuries in children (aged 0-4 yrs), (P), 2016/17	80.6	1/15	80.6	165.4	126.3	▼	●
2.07i - Hosp adm caused by unintentional and deliberate injuries in children (aged 0-14 yrs), (F), 2016/17	64.0	1/15	64.0	121.8	89.7	■	●
2.07i - Hosp adm caused by unintentional and deliberate injuries in children (aged 0-14 yrs), (M), 2016/17	85.0	3/15	73.5	130.7	112.7	■	●
2.07i - Hosp adm caused by unintentional and deliberate injuries in children (aged 0-14 yrs), (P), 2016/17	74.8	3/15	70.6	126.4	101.5	▼	●
2.07ii - Hosp adm caused by uninten and delib injuries in young people (aged 15-24 years), (F), 2016/17	113.0	3/15	111.2	233.5	126.3	■	●
2.07ii - Hosp adm caused by uninten and delib injuries in young people (aged 15-24 years), (M), 2016/17	109.9	5/15	98.4	165.8	131.8	■	●
2.07ii - Hosp adm caused by uninten and delib injuries in young people (aged 15-24 years), (P), 2016/17	111.4	3/15	104.7	195.8	129.2	▲	●
2.08i - Avg difficulties score for all LAC aged 5-16 who have been in care => 12 mths, (P), 2016/17	16.9	15/16	12.9	19.2	14.1	■	●
2.08ii - Percentage of children where there is a cause for concern, (P), 2016/17	50.0	14/16	31.4	59.2	38.1	■	●
2.09i - Smoking prevalence at age 15 - current smokers (WAY survey), (P), 2014/15	6.9	2/16	5.5	10.5	8.2	■	●
2.09ii - Smoking prevalence at age 15 - regular smokers (WAY survey), (P), 2014/15	4.5	2/16	3.2	7.2	5.5	■	●
2.09iii - Smoking prevalence at age 15 - occasional smokers (WAY survey), (P), 2014/15	2.5	7/16	2.1	4.7	2.7	■	●
2.10i - Emergency Hospital Admissions for Intentional Self-Harm, (F), 2016/17	214.3	4/15	203.0	386.3	233.7	■	●
2.10ii - Emergency Hospital Admissions for Intentional Self-Harm, (M), 2016/17	112.9	2/15	95.3	220.9	138.8	■	●
2.10ii - Emergency Hospital Admissions for Intentional Self-Harm, (P), 2016/17	161.8	4/15	151.2	297.6	185.3	■	●
2.11i - Proportion of the population meeting the recommended '5-a-day' on a 'usual day' (adults), (P), 2016/17	58.1	10/16	66.4	53.7	57.4	■	●
2.11ii - Average number of portions of fruit consumed daily (adults), (P), 2016/17	2.7	10/16	2.9	2.5	2.6	■	●
2.11iii - Average number of portions of vegetables consumed daily (adults), (P), 2016/17	2.6	15/16	2.9	2.6	2.7	■	●
2.11iv - Prop of the pop. meeting the recommended "5-a-day" at age 15, (P), 2014/15	54.4	7/16	60.3	48.5	52.4	■	●
2.11v - Average number of portions of fruit consumed daily at age 15 (WAY survey), (P), 2014/15	2.4	5/16	2.6	2.2	2.4	■	●
2.11vi - Average number of portions of vegetables consumed daily at age 15 (WAY survey), (P), 2014/15	2.5	3/16	2.7	2.3	2.4	■	●

**Statistical Significance compared to England or Benchmark:**

- Better
- Same
- Worse

**Direction of Travel:**

- ▲ Increasing
- ▼ Decreasing
- ▲ Increasing and getting better
- ▼ Decreasing and getting better
- ▲ Increasing and getting worse
- ▼ Decreasing and getting worse

**Other Indicators:**

- ▲ No significant change
- Cannot be calculated

Ranking, Best and Worst columns are compared to the nearest neighbours only. Rank: 1 is calculated as the best (or lowest when no polarity is applied)

Source: August 2018 PHOF, PHE

# Health Improvement (cont): Leicestershire

Indicator	Value	Rank	Best/Lowest	Worst/Highest	England	DoT	RAG
2.14 - Smoking Prevalence in adults - current smokers (APS), (F), 2017	12.0	5/16	9.9	15.5	13.0	—	●
2.14 - Smoking Prevalence in adults - current smokers (APS), (M), 2017	12.1	2/16	10.6	18.7	16.8	—	●
2.14 - Smoking Prevalence in adults - current smokers (APS), (P), 2017	25.7	2/16	23.1	34.2	31.8	—	●
2.15i - Successful completion of drug treatment - opiate users, (P), 2016	6.6	12/16	11.8	4.2	6.7	▼	●
2.15ii - Successful completion of drug treatment - non-opiate users, (P), 2016	35.7	8/16	48.4	21.8	37.1	▲	●
2.15iii - Successful completion of alcohol treatment, (P), 2016	38.8	10/16	51.8	27.0	38.7	▲	●
2.15iv - Deaths from drug misuse, (F), 2015 - 17	Null	Null	Null	Null	2.3	—	●
2.15iv - Deaths from drug misuse, (M), 2015 - 17	3.5	1/16	3.5	6.7	6.3	—	●
2.15iv - Deaths from drug misuse, (P), 2015 - 17	2.2	1/16	2.2	4.6	4.3	—	●
2.16 - Adults with SM treatment: need who engage in comm-based treatment after prison, (P), 2016/17	59.7	1/16	59.7	20.8	30.3	—	●
2.17 - Estimated diabetes diagnosis rate, (P), 2017	78.6	6/16	82.5	67.8	77.1	—	●
2.18 - Admission episodes for alcohol-related conditions - narrow definition, (F), 2016/17	471.6	5/15	356.8	616.6	473.2	—	●
2.18 - Admission episodes for alcohol-related conditions - narrow definition, (M), 2016/17	699.6	3/15	626.4	970.7	818.4	—	●
2.18 - Admission episodes for alcohol-related conditions - narrow definition, (P), 2016/17	577.6	4/15	489.0	766.2	636.4	—	●
2.19 - Cancer diagnosed at early stage (experimental statistics), (P), 2016	51.7	6/16	46.6	56.3	52.6	—	●
2.20i - Cancer screening coverage - breast cancer, (F), 2017	82.3	1/16	82.3	74.2	75.4	▼	●
2.20ii - Cancer screening coverage - cervical cancer, (F), 2017	77.4	4/16	78.7	71.4	72.0	▼	●
2.20iii - Cancer screening coverage - bowel cancer, (P), 2017	63.8	4/16	65.3	59.9	58.8	—	●
2.20iv - Abdominal Aortic Aneurysm Screening - Coverage, (M), 2016/17	81.1	15/16	87.8	80.1	80.9	—	●
2.22ii - Cumul % of the eligible pop aged 40-74 offered an NHS Health Check, (P), 2013/14 - 17/18	116.2	2/16	127.5	57.1	90.9	—	●
2.22iv - Cumul % of the eligible pop aged 40-74 offered and received an NHS Health Check, (P), 2013/14 - 17/18	43.1	13/16	58.5	31.9	48.7	—	●
2.22v - Cumul % of the eligible pop aged 40-74 who received an NHS Health Check, (P), 2013/14 - 17/18	50.1	4/16	59.0	28.8	44.3	—	●
2.23i - Self-reported wellbeing - people with a low satisfaction score, (P), 2016/17	Null	Null	Null	Null	4.5	—	●
2.23ii - Self-reported wellbeing - people with a low worthwhile score, (P), 2016/17	Null	Null	Null	Null	3.6	—	●
2.23iii - Self-reported wellbeing - people with a low happiness score, (P), 2016/17	8.2	9/16	5.9	10.1	8.5	—	●
2.23iv - Self-reported wellbeing - people with a high anxiety score, (P), 2016/17	19.9	12/16	14.1	21.0	19.9	—	●
2.24i - Emergency hospital admissions due to falls in people aged 65 and over, (F), 2016/17	1,898.7	1/15	1,898.7	2,948.7	2,395.6	—	●
2.24i - Emergency hospital admissions due to falls in people aged 65 and over, (M), 2016/17	1,237.0	1/15	1,237.0	1,944.8	1,714.9	—	●
2.24i - Emergency hospital admissions due to falls in people aged 65 and over, (P), 2016/17	1,607.3	1/15	1,607.3	2,514.5	2,113.8	—	●
2.24ii - Emergency hospital admissions due to falls in people aged 65-79, (F), 2016/17	941.2	6/15	880.5	1,406.3	1,152.5	—	●
2.24ii - Emergency hospital admissions due to falls in people aged 65-79, (M), 2016/17	553.4	1/15	553.4	887.8	813.8	—	●
2.24ii - Emergency hospital admissions due to falls in people aged 65-79, (P), 2016/17	756.1	4/15	744.6	1,156.8	993.3	—	●
2.24iii - Emergency hospital admissions due to falls in people aged 80+, (F), 2016/17	4,675.4	1/15	4,675.4	7,421.8	6,000.9	—	●
2.24iii - Emergency hospital admissions due to falls in people aged 80+, (M), 2016/17	3,219.4	1/15	3,219.4	5,010.0	4,328.4	—	●
2.24iii - Emergency hospital admissions due to falls in people aged 80+, (P), 2016/17	4,075.8	1/15	4,075.8	6,451.6	5,363.2	—	●

**Statistical Significance compared to England or Benchmark:**

- Better
- Same
- Worse

**Direction of Travel:**

- ▲ Increasing
- ▼ Decreasing
- ▲ Increasing and getting better
- ▼ Decreasing and getting better
- ▲ Increasing and getting worse
- ▼ Decreasing and getting worse

**Other Indicators:**

- ▲ No significant change
- Cannot be calculated

Ranking, Best and Worst columns are compared to the nearest neighbours only. Rank: 1 is calculated as the best (or lowest when no polarity is applied) Source: August 2018 PHOF, PHE

# Health Protection: Leicestershire

Indicator	Value	Rank	Best/Lowest	Worst/Highest	England	DoT	RAG
3.01 - Fraction of mortality attributable to particulate air pollution, (P), 2016	5.9	16/16	3.9	5.9	5.3	—	●
3.02 - Chlamydia detection rate (15-24 year olds), (F), 2017	2,515.0	4/16	3,125.1	1,513.9	2,501.8	▲	●
3.02 - Chlamydia detection rate (15-24 year olds), (M), 2017	1,341.2	3/16	1,495.8	734.4	1,263.6	▲	●
3.02 - Chlamydia detection rate (15-24 year olds), (P), 2017	1,886.5	4/16	2,285.2	1,217.3	1,881.9	▲	●
3.03i - Population vaccination coverage - Hepatitis B (1 year old), (P), 2016/17	100.0	4/13	100.0	0.0	Null	—	●
3.03i - Population vaccination coverage - Hepatitis B (2 years old), (P), 2016/17	64.3	10/12	100.0	0.0	Null	—	●
3.03ii - Population vaccination coverage - BCG - areas offering universal BCG only, (P), 2016/17	Null	Null	Null	Null	Null	—	●
3.03iii - Population vaccination coverage - Dtap / IPV / Hib (1 year old), (P), 2016/17	97.3	1/16	97.3	92.6	93.4	▲	●
3.03iii - Population vaccination coverage - Dtap / IPV / Hib (2 years old), (P), 2016/17	98.2	3/16	98.5	95.0	95.1	▲	●
3.03iv - Population vaccination coverage - MenC, (P), 2015/16	98.1	2/15	98.1	94.5	Null	—	●
3.03ix - Population vaccination coverage - MMR for one dose (5 years old), (P), 2016/17	98.3	3/16	98.7	93.0	95.0	▲	●
3.03v - Population vaccination coverage - PCV, (P), 2016/17	97.3	1/16	97.3	93.1	93.5	▲	●
3.03vi - Population vaccination coverage - Hib / MenC booster (5 years old), (P), 2016/17	96.4	3/16	96.9	89.4	92.6	▲	●
3.03vi - Population vaccination coverage - Hib / MenC booster (2 years old), (P), 2016/17	96.1	3/16	96.5	90.9	91.5	▲	●
3.03vii - Population vaccination coverage - PCV booster, (P), 2016/17	96.2	3/16	96.3	90.9	91.5	▲	●
3.03viii - Population vaccination coverage - MMR for one dose (2 years old), (P), 2016/17	96.2	2/16	97.2	91.1	91.6	▲	●
3.03x - Population vaccination coverage - MMR for two doses (5 years old), (P), 2016/17	93.8	2/16	95.9	85.1	87.6	▲	●
3.03xii - Pop vaccination coverage - HPV vaccination coverage for one dose (12-13 yrs), (F), 2016/17	93.5	4/16	97.9	83.3	87.2	—	●
3.03xiii - Population vaccination coverage - PPV, (P), 2016/17	72.4	7/16	75.7	65.6	69.8	▲	●
3.03xiv - Population vaccination coverage - Flu (aged 65+), (P), 2017/18	74.1	9/16	76.1	70.0	72.6	▲	●
3.03xv - Population vaccination coverage - Flu (at risk individuals), (P), 2017/18	48.5	12/16	53.4	45.3	48.9	▲	●
3.03xvi - Pop vaccination coverage - HPV vaccination coverage for two doses (13-14 yrs), (F), 2016/17	90.7	4/16	94.5	79.5	83.1	—	●
3.03xvii - Population vaccination coverage - Shingles vaccination coverage (70 years old), (P), 2016/17	55.4	2/16	56.0	43.2	48.3	—	●
3.03xviii - Population vaccination coverage - Flu (2-4 years old) - historical method, (P), 2016/17	49.3	3/16	50.2	39.8	38.1	—	●
3.03xviii - Population vaccination coverage - Flu (2-3 years old) - current method, (P), 2017/18	61.4	1/16	61.4	38.4	43.5	—	●
3.04 - HIV late diagnosis, (P), 2014 - 16	Null	Null	Null	Null	40.1	—	●
3.05i - Treatment completion for TB, (P), 2015	88.9	1/12	88.9	69.8	83.4	▲	●
3.05ii - Incidence of TB, (P), 2014 - 16	3.3	5/16	1.9	8.0	10.9	—	●
3.06 - NHS.org with a board app sustainable dev management plan, (N/A), 2015/16	40.0	15/16	83.3	28.6	66.2	▲	●
3.08 - Adjusted antibiotic prescribing in primary care by the NHS, (P), 2017	1.0	8/16	0.9	1.1	1.0	—	●

**Statistical Significance compared to England or Benchmark:**

- Better
- Same
- Worse
- Higher
- Lower
- Not compared

**Direction of Travel:**

- ▲ Decreasing
- ▲ Decreasing and getting better
- ▲ Decreasing and getting worse
- ▲ Increasing
- ▲ Increasing and getting better
- ▲ Increasing and getting worse
- ▲ No significant change
- Cannot be calculated

Ranking, Best and Worst columns are compared to the nearest neighbours only. Rank: 1 is calculated as the best (or lowest when no polarity is applied) Source: August 2018 PHOF, PHE



# Healthcare Premature Mortality: Leicestershire

Indicator	Value	Rank	Best/Lowest	Worst/Highest	England	DoT	RAG
4.01 - Infant mortality, (P), 2014 - 16	3.9	10/16	2.6	5.2	3.9	—	●
4.02 - Proportion of five year old children free from dental decay, (P), 2016/17	77.7	15/16	87.1	75.7	76.7	—	●
4.03 - Mortality rate from causes considered preventable, (F), 2014 - 16	119.7	7/16	103.6	146.4	138.5	—	●
4.03 - Mortality rate from causes considered preventable, (M), 2014 - 16	188.3	4/16	178.1	235.2	230.4	—	●
4.03 - Mortality rate from causes considered preventable, (P), 2014 - 16	152.8	4/16	139.0	189.5	182.8	—	●
4.04i - Under 75 mortality rate from all cardiovascular diseases, (F), 2014 - 16	39.6	10/16	30.6	45.1	45.8	—	●
4.04i - Under 75 mortality rate from all cardiovascular diseases, (M), 2014 - 16	86.8	6/16	74.1	106.1	102.7	—	●
4.04i - Under 75 mortality rate from all cardiovascular diseases, (P), 2014 - 16	62.8	8/16	51.7	75.0	73.5	—	●
4.04ii - Under 75 mortality rate from cardiovascular diseases considered preventable, (F), 2014 - 16	20.9	10/16	14.4	25.3	24.3	—	●
4.04ii - Under 75 mortality rate from cardiovascular diseases considered preventable, (M), 2014 - 16	60.8	9/16	51.1	73.1	70.4	—	●
4.04ii - Under 75 mortality rate from cardiovascular diseases considered preventable, (P), 2014 - 16	40.5	9/16	32.2	48.8	46.7	—	●
4.05i - Under 75 mortality rate from cancer, (F), 2014 - 16	116.8	9/16	104.8	130.8	122.6	—	●
4.05i - Under 75 mortality rate from cancer, (M), 2014 - 16	130.5	3/16	125.8	157.9	152.1	—	●
4.05i - Under 75 mortality rate from cancer, (P), 2014 - 16	123.4	5/16	114.7	142.8	136.8	—	●
4.05ii - Under 75 mortality rate from cancer considered preventable, (F), 2014 - 16	66.1	7/16	58.2	80.2	73.4	—	●
4.05ii - Under 75 mortality rate from cancer considered preventable, (M), 2014 - 16	70.0	3/16	64.2	89.7	85.9	—	●
4.05ii - Under 75 mortality rate from cancer considered preventable, (P), 2014 - 16	68.0	6/16	61.5	84.8	79.4	—	●
4.06i - Under 75 mortality rate from liver disease, (F), 2014 - 16	10.7	9/16	8.7	15.2	12.8	—	●
4.06i - Under 75 mortality rate from liver disease, (M), 2014 - 16	16.1	4/16	15.3	25.3	23.9	—	●
4.06i - Under 75 mortality rate from liver disease, (P), 2014 - 16	13.3	6/16	11.9	19.1	18.3	—	●
4.06ii - Under 75 mortality rate from liver disease considered preventable, (F), 2014 - 16	9.0	9/16	7.0	13.1	10.9	—	●
4.06ii - Under 75 mortality rate from liver disease considered preventable, (M), 2014 - 16	14.9	6/16	13.6	23.0	21.5	—	●
4.06ii - Under 75 mortality rate from liver disease considered preventable, (P), 2014 - 16	11.9	7/16	10.4	17.2	16.1	—	●
4.07i - Under 75 mortality rate from respiratory disease, (F), 2014 - 16	20.2	2/16	19.1	29.0	28.7	—	●
4.07i - Under 75 mortality rate from respiratory disease, (M), 2014 - 16	29.9	4/16	26.3	41.1	39.2	—	●
4.07i - Under 75 mortality rate from respiratory disease, (P), 2014 - 16	24.9	5/16	23.3	34.6	33.8	—	●
4.07ii - Under 75 mortality rate from respiratory disease considered preventable, (F), 2014 - 16	10.4	1/16	10.4	17.1	16.5	—	●
4.07ii - Under 75 mortality rate from respiratory disease considered preventable, (M), 2014 - 16	12.9	2/16	11.7	23.0	20.8	—	●
4.07ii - Under 75 mortality rate from respiratory disease considered preventable, (P), 2014 - 16	11.6	2/16	11.1	19.7	18.6	—	●
4.08 - Mortality rate from a range of specified communicable diseases, incl influenza, (F), 2014 - 16	5.5	1/16	5.5	12.2	9.9	—	●
4.08 - Mortality rate from a range of specified communicable diseases, incl influenza, (M), 2014 - 16	6.1	1/16	6.1	14.7	11.6	—	●
4.08 - Mortality rate from a range of specified communicable diseases, incl influenza, (P), 2014 - 16	5.9	1/16	5.9	13.3	10.7	—	●
4.09i - Excess under 75 mortality rate in adults with serious mental illness, (P), 2014/15	362.9	11/16	294.5	414.9	370.0	—	●
4.09ii - Proportion of adults in the pop in contact with secondary mental health services, (P), 2014/15	4.3	6/16	3.7	6.7	5.4	—	●

**Statistical Significance compared to England or Benchmark:**

- Better
- Same
- Worse

**Direction of Travel:**

- ▲ Increasing
- ▼ Decreasing
- ▲ Increasing and getting better
- ▼ Decreasing and getting better
- ▲ Increasing and getting worse
- ▼ Decreasing and getting worse

▲ No significant change  
■ Cannot be calculated

Ranking, Best and Worst columns are compared to the nearest neighbours only. Rank: 1 is calculated as the best (or lowest when no polarity is applied) Source: August 2018 PHOF, PHE

# Healthcare Premature Mortality (cont): Leicestershire

Indicator	Value	Rank	Best/Lowest	Worst/Highest	England	DoT	RAG
4.10 - Suicide rate, (F), 2014 - 16	4.3	6/16	3.8	6.0	4.8	—	●
4.10 - Suicide rate, (M), 2014 - 16	14.1	5/16	12.4	19.0	15.3	—	●
4.10 - Suicide rate, (P), 2014 - 16	8.9	5/16	8.2	12.2	9.9	—	●
4.11 - Emergency readmissions within 30 days of discharge from hospital, (F), 2011/12	11.4	14/16	10.1	11.7	11.5	—	●
4.11 - Emergency readmissions within 30 days of discharge from hospital, (M), 2011/12	11.6	11/16	10.6	12.6	12.1	—	●
4.11 - Emergency readmissions within 30 days of discharge from hospital, (P), 2011/12	11.5	14/16	10.3	12.1	11.8	—	●
4.12i - Preventable sight loss - age related macular degeneration (AMD), (P), 2016/17	110.6	12/16	35.0	158.4	111.3	▲	●
4.12ii - Preventable sight loss - glaucoma, (P), 2016/17	9.3	9/16	5.6	21.0	13.1	▲	●
4.12iii - Preventable sight loss - diabetic eye disease, (P), 2016/17	2.4	5/15	1.0	4.8	3.1	▲	●
4.12iv - Preventable sight loss - sight loss certifications, (P), 2016/17	48.8	12/16	17.1	75.4	42.4	▲	●
4.13 - Health related quality of life for older people, (P), 2016/17	0.8	10/16	0.8	0.7	0.7	—	●
4.14i - Hip fractures in people aged 65 and over, (F), 2016/17	751.0	15/15	600.3	751.0	693.2	—	●
4.14i - Hip fractures in people aged 65 and over, (M), 2016/17	376.1	4/15	351.6	429.9	407.9	—	●
4.14i - Hip fractures in people aged 65 and over, (P), 2016/17	598.8	15/15	499.2	598.8	575.0	—	●
4.14ii - Hip fractures in people aged 65 and over - aged 65-79, (F), 2016/17	322.5	13/15	237.7	349.6	306.3	—	●
4.14ii - Hip fractures in people aged 65 and over - aged 65-79, (M), 2016/17	159.8	11/15	122.2	171.6	166.3	—	●
4.14ii - Hip fractures in people aged 65 and over - aged 65-79, (P), 2016/17	246.0	13/15	182.9	258.4	240.6	—	●
4.14iii - Hip fractures in people aged 65 and over - aged 80+, (F), 2016/17	1,993.9	15/15	1,615.4	1,993.9	1,815.2	—	●
4.14iii - Hip fractures in people aged 65 and over - aged 80+, (M), 2016/17	1,003.5	3/15	961.5	1,277.6	1,108.4	—	●
4.14iii - Hip fractures in people aged 65 and over - aged 80+, (P), 2016/17	1,621.9	13/15	1,364.4	1,629.2	1,544.5	—	●
4.15i - Excess winter deaths index (single year, all ages), (F), Aug 2015 - Jul 2016	9.1	1/16	9.1	21.6	16.2	—	●
4.15i - Excess winter deaths index (single year, all ages), (M), Aug 2015 - Jul 2016	14.7	12/16	5.1	19.4	13.9	—	●
4.15i - Excess winter deaths index (single year, all ages), (P), Aug 2015 - Jul 2016	11.8	3/16	8.7	19.3	15.1	—	●
4.15ii - Excess winter deaths index (single year, age 85+), (F), Aug 2015 - Jul 2016	8.2	2/16	7.1	25.8	17.8	—	●
4.15ii - Excess winter deaths index (single year, age 85+), (M), Aug 2015 - Jul 2016	22.5	14/16	6.6	30.5	17.5	—	●
4.15ii - Excess winter deaths index (single year, age 85+), (P), Aug 2015 - Jul 2016	13.5	6/16	11.8	22.6	17.7	—	●
4.15iii - Excess winter deaths index (3 years, all ages), (F), Aug 2013 - Jul 2016	15.7	1/16	15.7	24.7	20.2	—	●
4.15iii - Excess winter deaths index (3 years, all ages), (M), Aug 2013 - Jul 2016	16.0	7/16	9.7	18.8	15.4	—	●
4.15iii - Excess winter deaths index (3 years, all ages), (P), Aug 2013 - Jul 2016	15.8	3/16	14.7	20.8	17.9	—	●
4.15iv - Excess winter deaths index (3 years, age 85+), (F), Aug 2013 - Jul 2016	21.9	4/16	20.5	29.0	25.3	—	●
4.15iv - Excess winter deaths index (3 years, age 85+), (M), Aug 2013 - Jul 2016	28.1	13/16	18.6	34.1	23.3	—	●
4.15iv - Excess winter deaths index (3 years, age 85+), (P), Aug 2013 - Jul 2016	24.2	10/16	20.7	30.9	24.6	—	●
4.16 - Estimated dementia diagnosis rate (aged 65+), (P), 2018	70.0	3/16	75.4	59.4	67.5	—	●

**Statistical Significance compared to England or Benchmark:**

- Better
- Same
- Worse
- Higher
- Lower
- Not compared

**Direction of Travel:**

- ▲ Increasing
- ▼ Decreasing
- ▲ Increasing and getting better
- ▼ Decreasing and getting better
- ▲ Increasing and getting worse
- ▼ Decreasing and getting worse

▲ No significant change  
 ▼ Cannot be calculated

Ranking, Best and Worst columns are compared to the nearest neighbours only. Rank: 1 is calculated as the best (or lowest when no polarity is applied) Source: August 2018 PHOF, PHE

If you require information contained in this leaflet in another version e.g. large print, Braille, tape or alternative language please telephone: 0116 305 6803, Fax: 0116 305 7271 or Minicom: 0116 305 6160.

જો આપ આ માહિતી આપની ભાષામાં સમજવામાં થોડી મદદ ઇચ્છતાં હો તો 0116 305 6803 નંબર પર ફોન કરશો અને અમે આપને મદદ કરવા યત્ન કરીશું.

ਜੇਕਰ ਤੁਹਾਨੂੰ ਇਸ ਜਾਣਕਾਰੀ ਨੂੰ ਸਮਝਣ ਵਿਚ ਕੁਝ ਮਦਦ ਚਾਹੀਦੀ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ 0116 305 6803 ਨੰਬਰ ਤੇ ਫੋਨ ਕਰੋ ਅਤੇ ਅਸੀਂ ਤੁਹਾਡੀ ਮਦਦ ਲਈ ਕਿਸੇ ਦਾ ਪ੍ਰਬੰਧ ਕਰ ਦਵਾਂਗੇ।

এই তথ্য নিজের ভাষায় বুঝার জন্য আপনার যদি কোন সাহায্যের প্রয়োজন হয়, তবে 0116 305 6803 এই নম্বরে ফোন করলে আমরা উপযুক্ত ব্যক্তির ব্যবস্থা করবো।

اگر آپ کو یہ معلومات سمجھنے میں کچھ مدد درکار ہے تو براہ مہربانی اس نمبر پر کال کریں اور ہم آپ کی مدد کے لئے کسی کا انتظام کر دیں گے۔ 0116 305 6803

假如閣下需要幫助，用你的語言去明白這些資訊，請致電 0116 305 6803，我們會安排有關人員為你提供幫助。

Jeżeli potrzebujesz pomocy w zrozumieniu tej informacji w Twoim języku, zadzwoń pod numer 0116 305 6803, a my Ci dopomożemy.

Strategic Business Intelligence Team  
Strategy and Business Intelligence Branch  
Chief Executive's Department  
Leicestershire County Council  
County Hall  
Glenfield  
Leicester  
LE3 8RA  
[ri@leics.gov.uk](mailto:ri@leics.gov.uk)  
[www.lsr-online.org](http://www.lsr-online.org)